

S3 Table. Review of SFRP1 in the hair follicle.

Model	SFRP1 mRNA/protein	SFRP1 location	SFRP1 biological effect?	Changes in β-catenin signalling by SFRP1?	Ref
Human frozen skin	mRNA	Melanocyte precursors within bulge region	n/a	x	1
Murine	mRNA	Anagen and telogen HFSCs	n/a	x	2
Murine; K14-H2BGFP Lef1-RFP Sox2GFP K14-RFP Sox9-GFP Crabp1-GFP	mRNA	ORS, Mx, DP, HFSCs	n/a	x	3
C57BL/6 skin	mRNA	n/a	n/a	x	4
C57BL/6 neonatal	mRNA	Dermal cells	Impaired HF formation	x	5
BALB/C mice	mRNA	n/a	n/a	x	6
Human isolated DP	mRNA	DP	n/a	x	7
HrHp/HrHp mice	mRNA	n/a	n/a	x	8
HrHp/HrHp mice	mRNA	n/a	n/a	x	9
Sox2eGFP mice	mRNA	Dermal papilla populations GFP-CD133- GFP-CD133+ GFP+ CD133+	n/a	x	10
Human frozen skin	mRNA	HFSCs	n/a	x	11
Immortalized balding and non-balding DP cells	mRNA	DP cells	n/a	x	12

1. Goldstein NB, Koster MI, Hoaglin LG, et al. Isolating RNA from precursor and mature melanocytes from human vitiligo and normal skin using laser capture microdissection. *Exp Dermatol.* 2016;25(10):805-811.
2. Lim X, Tan SH, Yu K Lou, Lim SBH, Nusse R. Axin2 marks quiescent hair follicle bulge stem cells that are maintained by autocrine Wnt/β-catenin signaling. *Proc Natl Acad Sci.* 2016;113(11):E1498-E1505.
3. Rezza A, Wang Z, Sennett R, et al. Signaling networks among stem cell precursors, transit-amplifying progenitors, and their niche in developing hair follicles. *Cell Rep.* 2016;14(12):3001-3018.

4. Lee J, Lee K, Chung H. Investigation of transcriptional gene profiling in normal murine hair follicular substructures using next-generation sequencing to provide potential insights into skin disease. *Cell Transpl.* 2016;25(2):377-399.
5. Bak S-S, Kim MK, Kim JC, Sung YK. Follistatin and secreted frizzled-related protein 1, OVO homolog-like 1-regulated genes, are important for hair follicle neogenesis. *Exp Dermatol.* 2015;24(7):550-551.
6. Kim BK, Yoon SK. Expression of Sfrp2 is increased in catagen of hair follicles and inhibits keratinocyte proliferation. *Ann Dermatol.* 2014;26(1):79-87.
7. Higgins CA, Chen JC, Cerise JE, Jahoda CAB, Christiano AM. Microenvironmental reprogramming by three-dimensional culture enables dermal papilla cells to induce de novo human hair-follicle growth. *Proc Natl Acad Sci U S A.* 2013;110(49):19679-19688.
8. Choi JH, Kim BK, Kim JK, Lee HY, Park J, Yoon SK. Downregulation of Foxe1 by HR suppresses Msx1 expression in the hair follicles of Hr HP mice. *BMB Rep.* 2011;44(7):478-483.
9. Kim B-K, Baek I-C, Lee H-Y, Kim J-K, Song H-H, Yoon SK. Gene expression profile of the skin in the “hairpoor” (HrHp) mice by microarray analysis. *BMC Genomics.* 2010;11:640.
10. Driskell RR, Giangreco A, Jensen KB, Mulder KW, Watt FM. Sox2-positive dermal papilla cells specify hair follicle type in mammalian epidermis. *Development.* 2009;136(16):2815-2823.
11. Ohyama M, Terunuma A, Tock CL, et al. Characterization and isolation of stem cell-enriched human hair follicle bulge cells. *J Clin Invest.* 2006;116(1):249-260.
12. Chew EGY, Tan JHJ, Bahta AW, et al. Differential expression between human dermal papilla cells from balding and non-balding scalps reveals new candidate genes for androgenetic alopecia. *J Invest Dermatol.* 2016;136(8):1559-1567.